

Appendix D

Kern Water Bank Raptor and Upland Bird Survey

Report: August 2012 - May 2016



Mourning Dove (*Zenaida macroura*)

Sterling Wildlife Biology

Woodland CA 95695
Phone: 530 908-3836
E-Mail: jsterling@wavecable.com
Web: www.sterlingbirds.com

Kern Water Bank

Raptor and Upland Bird Survey Report: August 2012 - May 2016

23 May 2016

Introduction

The property managed by the Kern Water Bank Authority supports a wealth of native wildlife, especially an abundance of upland birds and raptors attracted to the recharge ponds and/or the upland habitats. In order to document and quantify this natural resource value, John Sterling of Sterling Wildlife Biology conducted bird surveys from late August 2012 to late May 2016. These surveys are intended to capture a snapshot of the bird use of the project area during the winter, spring/fall migration and the breeding seasons. The resulting data serve to document the regional importance of habitats on the Kern Water Bank for raptors and upland birds during this period. Most importantly, the data describe the baseline of existing conditions that may be used to inform range management practices with regard to productive bird habitat. This baseline data will be used to measure population trends with range management enhancement and/or unmanaged changes in habitat due to the extension or end of current drought conditions.

Methods

Survey Methods

For the raptor/Loggerhead Shrike and upland bird surveys, John Sterling visited the sites approximately every two weeks for a total of eighty-eight surveys. The dates of the surveys were approximately every two weeks starting on 31 August 2012 to 16-17 May 2016, with breaks in June and July. Raptor/Loggerhead Shrike surveys were conducted in June and July only in 2015. Upland bird surveys were not conducted during the summer period as most nesting had been completed by 31 May and there were few birds remaining on the study area until fall migration began in mid August. Upland bird surveys were conducted on fixed, one-half mile long transects (Figure 1). Mr. Sterling conducted upland bird surveys by walking transects and recording all birds heard or seen within 200 meters of the transect line. He tabulated the numbers of each species. Each transect was surveyed up to eighty-two times with the exception of new transects that were created in October 2012 and surveyed eighty-one times. Transects were 0.5 miles long with the exception of Transect G, which was 0.25 miles long due to the small size of that habitat fragment. For sixty-three sets of raptor surveys, Mr. Sterling drove most roads to cover the entire project area and kept running tallies of numbers of individuals of all raptor species and Loggerhead Shrike. All data were compiled onto spreadsheets (See attached Appendix B & C files).

Descriptions of Upland Bird Survey Transects

The following are brief descriptions of the bird habitat along each of the survey transects including photographs showing conditions on 7 June 2013.

Transect A
The transect borders a large canal that is watered and supports a few water birds. As such, it also supports tules and some sunflowers and other ruderal plants along its edge. There are several large willow trees (*Salix sp.*) but the habitat is mostly open, ruderal fields with some tumbleweed cover (*Salsola sp.*). During wet years, the ruderal vegetation is rank and relatively tall (up to 4 ft).



Figure 1. Locations of Upland Bird Survey Transects on the Kern Water Bank



Transect B

This transect borders a canal that was watered until spring 2012. It supports several willow trees along its banks along with mulefat, thistles and other ruderal vegetation. The fields are dry ponds and support ruderal vegetation.



Transect C

This transect is a honey mesquite (*Prosopis glandulosa*) woodland with some tree tobacco, annual grasses and some ruderal vegetation.



Transect D
The west side of this transect is a dry pond that is now an open willow woodland with moderate ruderal and annual grassland cover. The east side is a dry pond that is now a ruderal field with low, sparse vegetative cover.



Transect E
This transect has a honey mesquite woodland on the south side, with some annual grasses, but otherwise little vegetative cover apart from the mesquite. On the north side is a dry pond that is a ruderal field.



Transect F

This transect is relatively barren with some grasses, forbs and in some years dominated by tumbleweed.



Transect G

This transect has several honey mesquite shrubs on the east side, but the west side is dominated by saltbush (*Atriplex sp.*).



Transect H
This transect has some Fremont cottonwood saplings, along with an open honey mesquite woodland and tall ruderal vegetation on the west side. The east side is a dry pond and now a ruderal field.



Transect I
This transect has two small willow trees in a field dominated by tumbleweed on the south side, while the north side is an alfalfa field on property adjacent to the project area.



Special-Status Species Criteria

In evaluating the potential presence of special-status species, the following criteria were used to determine which species should be included:

- Bird species listed, or proposed for listing, as threatened or endangered under the ESA (50 CFR 17.11 [listed animals], and various notices in the Federal Register [proposed species]);
- bird species that are candidates for possible future listing as threatened or endangered under the ESA (61 FR 40: 7596-7613, February 28, 1996);
- bird species listed, or proposed for listing, by the State of California as threatened or endangered under CESA (14 CCR 670.5);
- bird species that meet the definitions of rare or endangered under CEQA (CEQA Guidelines, Section 15380);
- bird species of special concern to CDFG (CDFG in preparation [birds, Shuford and Gardali 2008];
- bird species fully protected in California (California Fish and Game Code, Section 3511 [birds]; and
- bird species included in CDFG's list of special animals and monitored by the California Natural Diversity Database (CNDDDB).

Results

Two hundred and nine species of birds have been recorded thus far at the Kern Water Bank during water bird, upland bird and raptor surveys since this project began in mid October 2011 (Appendix A). Many of those are discussed below or in the previous reports (Sterling Wildlife Biology, 27 April 2012, 9 December 2013, and 1 June 2015).

Upland Birds

One hundred and twelve species of birds were detected during the upland bird surveys. Of the nine transects, Transects A and C have the largest number of species with seventy-eight and seventy-seven, respectively (Figure 2). Although species richness (number of species) did not vary greatly over time in each transect, numbers of birds counted fluctuated greatly (Figures 3-11). Transects with the most birds contained mesquite and/or willow trees although Transect I with its grassland and alfalfa harbored large numbers of sparrows during the winter.

Figure 2. Cumulative Number of Species Found in each Transect




Figure 3. Number of Birds and Bird Species: Transect A.




Figure 4. Number of Birds and Bird Species: Transect B.




Figure 5. Number of Birds and Bird Species: Transect C.




Figure 6. Number of Birds and Bird Species: Transect D.




Figure 7. Number of Birds and Bird Species: Transect E.




Figure 8. Number of Birds and Bird Species: Transect F.




Figure 9. Number of Birds and Bird Species: Transect G.




Figure 10. Number of Birds and Bird Species: Transect H.





Figure 11. Number of Birds and Bird Species: Transect I.



Raptors and Shrikes

The comprehensive survey for raptors and Loggerhead Shrikes on the entire project area resulted in high numbers of raptors including Red-tailed Hawks and Loggerhead Shrikes (Figures 20-22), but also documented fourteen species of raptors using upland habitats during the surveys (Appendix B). Overall numbers of raptors dipped sharply after the winter of 2012-2013 and

then steadily decline to fewer than twenty individuals from February 2014 through May 2015, then consistently over twenty from October 2015 to March 2016. Conversely, Loggerhead Shrikes rebounded during the breeding season in 2015 after a similar decline (Figure 21). The increase from ten to fifty-five during a two-month period in spring 2015 was due to good reproductive success of local breeding population. The primary difference among the habitat conditions between spring of 2014 and 2015 was the lack of grasses and forbs in 2014 that resulted in low prey populations (large insects and lizards) in contrast to the tremendous amount of grasses and forbs in winter and spring of 2015. Although the amount of grasses and forbs were lower in 2016, the higher population maintained through the winter of 2015-2016 led to a higher breeding population that also had good reproductive success.

Figure 12. Raptor Population: 2012- 2016





Figure 13. Loggerhead Shrike Population: 2012-2016



Special Status Bird Species


There have been twenty-five special-status bird species found during the raptor and upland bird surveys since the project began in October 2011 (Table 2) with an additional fourteen species of waterbirds found during waterbird surveys in winter 2011-12.

Table 1. Species Status Bird Species (Landbirds and Raptors) found on the Kern Water Bank

Species Name	Conservation/Legal Status	Seasonal Status	Habitat	Record Dates
Cooper's Hawk	CA Watch List	Migration, Winter, Potential nesting	Nests in trees, hunts in woodlands and open grasslands	Daily during migration
White-tailed Kite	State Fully Protected	Migration, Winter, Potential nesting	Nests in trees, hunts in open grasslands	Daily during migration and winter in 2011-2013, but absent since then
Northern Harrier	CA Species of Special Concern	Migration, Winter, Potential nesting	Nests on the ground, hunts in wetlands and open grasslands	Daily during migration and winter in 2011-2013, but mostly absent since then with a few scattered records of individual migrants
Swainson's Hawk	CA Threatened Species	Nesting, Migration, Winter	Nests in trees, hunts in open grasslands	Nesting in summer 2012, scattered winter records in 2011-12; up to five individuals in Mar-May 2013; up to three individuals in Mar-May 2014; and up to four individuals in Apr/May 2015. No nest located on water bank property in 2015. Three active nests on water bank property in 2016 (Figure 14). Approximate locations: 1) 35°20'35.59"N, 119°20'27.20"W; 2) 35°20'43.52"N, 119°15'42.37"W; and 3) 35°19'11.17"N, 119°13'15.58"W
Ferruginous Hawk	CA Watch List	Winter	Hunts in open grasslands	Daily during winter, rare dark morph individual in March 2016
Golden Eagle	State Fully Protected and Federal Eagle Protection Act	Winter	Hunts in open grasslands	Four winter records
Osprey	CA Watch List	Migration, Winter	Hunts in wetlands and canals	Daily during winter 2011-12, only a few sightings of migrants since then
Mountain Plover	CA Species of Special Concern and Federal Proposed Threatened	Migration, Winter	Roosts and forages in grasslands	Two on 14 October 2013
Long-billed Curlew	CA Watch List and Federal Bird of Conservation Concern	Migration, Winter	Roosts and forages in grasslands and wetlands	Scattered winter and migration records
Burrowing Owl	CA Species of Special Concern	Nesting, Migration, Winter	Nests and hunts in grasslands	Found on most visits through year, and nested each spring
Vaux's Swift	CA Species of Special Concern	Migration	Forages over wetlands and grasslands	Found during spring migration visits
Merlin	CA Watch List and Federal Bird of		Hunts in grasslands and wetlands	Regular during late fall and winter

	Conservation Concern			
Peregrine Falcon	Federal Bird of Conservation Concern	Migration, Winter	Hunts in grasslands and wetlands	Regular during fall and winter of 2011-12, scattered records since then
Prairie Falcon	CA Watch List and Federal Bird of Conservation Concern	Migration, Winter	Hunts in grasslands	Found on most visits from Nov through Mar
Nuttall's Woodpecker	Federal Bird of Conservation Concern	Nesting, Migration, Winter	Nests in trees, forages in woodlands	Found during most survey visits on Transect C.
Willow Flycatcher	CA Endangered Species	Migration	Roosts in trees, hunts in open woodlands forages	Regular during migration
Vermilion Flycatcher	CA Species of Special Concern	Migration, Winter. Potential nesting	Nests in trees, forages in open woodlands and scrublands	Several winter records of at least three individuals since 2011
Loggerhead Shrike	CA Species of Special Concern and Federal Bird of Conservation Concern	Nesting, Migration, Winter	Nests in trees, hunts in open woodlands and scrublands	Found during each survey visit.
California Horned Lark	CA Watch List	Nesting, Migration, Winter	Nests on ground, forages in barren fields with little grassland cover	Found during each survey visit. Many breeding during 2015 and 2016.
Purple Martin	CA Species of Special Concern	Migration	Forages over wetlands and grasslands	1 Apr 2012
Lucy's Warbler	CA Species of Special Concern	Migration	Nests in trees, hunts in open woodlands and scrublands	1-4 Oct 2012; second record for the entire Central Valley
Grasshopper Sparrow	CA Species of Special Concern	Migration, Winter, Potential nesting	Nests on ground in grasslands	13 Nov, 10 Dec 2013, 23 Oct 2015
Tricolored Blackbird	CA Species of Special Concern and Federal Bird of Conservation Concern	Nesting, Migration, Winter	Nests in ruderal and marsh vegetation, forages in grasslands, fields and wetlands	Nesting in summer 2012, found most days in migration and winter during 2011-2012; nesting off site in 2015 but foraging on the water bank property
Yellow-headed Blackbird	CA Species of Special Concern	Migration, Winter. Potential nesting	Nests in marsh vegetation, forages in grasslands, fields and wetlands	Regular during migration and winter in 2011-12
Lawrence's Goldfinch	Federal Bird of Conservation Concern	Migration, Winter, Potential nesting	Nests in trees, forages in open woodlands and scrublands	Two late fall records in 2013, pairs on 23 Apr 2015, 1 Apr and 9 May 2016. Flocks Oct 2015-Jan 2016.

Figure 14. Active Swainson's Hawk nest locations in 2016.



Rare Birds

A few birds were discovered during the surveys that are not special-status species, but out of their normal range. These records are very important to our understanding of vagrancy in birds and the data are archived by county editors for "North American Birds" magazine and the online eBird database (administered by Cornell University's Laboratory of Ornithology). During fall migration two Black-throated Sparrows were found on Transects A and C. This desert species is very rare in the Central Valley. A fall migrant Clay-colored Sparrow was in mesquite and cottonwoods between transects A and B. This midwestern species is rare anywhere in California and especially in the Central Valley from which there are fewer than ten documented records. Surprisingly, no fewer than eight Brewer's Sparrows were found wintering in 2012-13 and several have been found each winter thereby establishing the area as a regular wintering area. Before the project there were very few documented records of this Great Basin and desert species during winter months in the Central Valley. There have been seven records of migrant Sage Thrashers—a Great Basin species, which is a rare but annual migrant in the Central Valley. During a 2012 fall survey, a Chestnut-collared Longspur was heard calling in flight over Transect I. This is a very rare wintering bird in the San Joaquin Valley and Tulare Basin with fewer than ten records. On a Christmas Bird Count before these surveys began, an Eastern Phoebe was documented for one of very few San Joaquin Valley and Tulare Basin records of this eastern species, which rarely occurs in California. During a spring surveys on 1 April 2012 and 12 March 2015, single Cassin's Kingbirds were found establishing the only Tulare Basin records away from eastern Bakersfield (only one record from nearby Kings County). Also on 1 April 2012, a male Purple Martin was photographed migrating over grasslands for one of few records for the Tulare Basin and San Joaquin Valley. Two rare warblers, Lucy's and Virginia's were found during fall migration for only the second and third records for the Central Valley, respectively. The Kern Water Bank has exceptional habitats for birds and many rare birds will likely be found and documented in the future dependent upon survey efforts.

Discussion

The bird use of property managed by the Kern Water Bank Authority is clearly very high in accordance to the large acreages of upland habitats. Overall, in terms of bird abundance, species diversity, acreage, location and habitat diversity, it is an important area of upland habitat, especially when compared to surrounding agricultural lands. These surveys documented particularly large populations raptors and shrikes, sparrows, and many other species typical of native upland habitats on the San Joaquin Valley floor. Of particular interest were the differences in the effect of the drought conditions among the years. There was measureable precipitation in winter of 2014-2015 and in the spring of 2015 with lesser amounts in winter 2015-2016 and spring 2016, which resulted in much growth of grasses and forbs throughout the water bank property. This was in contrast

to no new growth during spring 2014 that left the area devoid of grasses and forbs. As a result, Loggerhead Shrike populations rebounded to pre-winter 2013-2014 levels, primarily as a result of good reproductive success of local breeders. These shrikes prey upon large insects and lizards that were common during the springs of 2015 and 2016. In contrast, raptor counts remained low compared to wet years likely due to continuing low population levels of rodent prey, but did slightly increase in winter 2015-2016 through spring 2016 in response to relief of severe drought conditions.

Appendix A. List of Bird Species Recorded at the Kern Water Bank

Compiled By John Sterling (22 May 2016)

Bold-faced names = species rare in the Tulare Basin

Anseriformes - Screamers, Swans, Geese, and Ducks

Anatidae - Ducks, Geese, and Swans

Greater White-fronted Goose *Anser albifrons*

Snow Goose *Chen caerulescens*

Ross's Goose *Chen rossii*

Cackling Goose *Branta hutchinsii*

Canada Goose *Branta canadensis*

Tundra Swan *Cygnus columbianus*

Wood Duck *Aix sponsa*

Gadwall *Anas strepera*

Eurasian Wigeon *Anas penelope*

American Wigeon *Anas americana*

Mallard *Anas platyrhynchos*

Blue-winged Teal *Anas discors*

Cinnamon Teal *Anas cyanoptera*

Northern Shoveler *Anas clypeata*

Northern Pintail *Anas acuta*

Green-winged Teal *Anas crecca*

Canvasback *Aythya valisineria*

Redhead *Aythya americana*

Ring-necked Duck *Aythya collaris*

Greater Scaup *Aythya marila*

Lesser Scaup *Aythya affinis*

Bufflehead *Bucephala albeola*

Common Goldeneye *Bucephala clangula*

Barrow's Goldeneye *Bucephala islandica*

Hooded Merganser *Lophodytes cucullatus*

Common Merganser *Mergus merganser*

Red-breasted Merganser *Mergus serrator*

Ruddy Duck *Oxyura jamaicensis*

Galliformes - Gallinaceous Birds

Odontophoridae - New World Quail

California Quail *Callipepla californica*

Phasianidae - Partridges, Grouse, Turkeys, and Old World Quail

Ring-necked Pheasant *Phasianus colchicus* - I

Podicipediformes - Grebes

Podicipedidae - Grebes

Pied-billed Grebe *Podilymbus podiceps*

Horned Grebe *Podiceps auritus*

Eared Grebe *Podiceps nigricollis*

Western Grebe *Aechmophorus occidentalis*

Clark's Grebe *Aechmophorus clarkii*

Phalacrocoracidae - Cormorants

Double-crested Cormorant *Phalacrocorax auritus*

Pelecaniformes - Pelicans, Herons, Ibises, and Allies

Pelecanidae - Pelicans

American White Pelican *Pelecanus erythrorhynchos*

Ardeidae - Herons, Bitterns, and Allies

Great Blue Heron *Ardea herodias*

Great Egret *Ardea alba*

Snowy Egret *Egretta thula*

Cattle Egret *Bubulcus ibis*

Green Heron *Butorides virescens*

Black-crowned Night-Heron *Nycticorax nycticorax*

Threskiornithidae - Ibises and Spoonbills

White-faced Ibis *Plegadis chihi*

Accipitriformes - Hawks, Kites, Eagles, and Allies

Cathartidae - New World Vultures

Turkey Vulture *Cathartes aura*

Pandionidae - Ospreys

Osprey *Pandion haliaetus*

Accipitridae - Hawks, Kites, Eagles, and Allies

White-tailed Kite *Elanus leucurus*

Bald Eagle *Haliaeetus leucocephalus*

Northern Harrier *Circus cyaneus*

Sharp-shinned Hawk *Accipiter striatus*

Cooper's Hawk *Accipiter cooperii*

Red-shouldered Hawk *Buteo lineatus*

Swainson's Hawk *Buteo swainsoni*

Red-tailed Hawk *Buteo jamaicensis*

Ferruginous Hawk *Buteo regalis*

Golden Eagle *Aquila chrysaetos*

Gruiformes - Rails, Cranes, and Allies

Rallidae - Rails, Gallinules, and Coots

Virginia Rail *Rallus limicola*

Sora *Porzana carolina*

Common Gallinule *Gallinula galeata*

American Coot *Fulica americana*

Charadriiformes - Shorebirds, Gulls, Auks, and Allies

Recurvirostridae - Stilts and Avocets

Black-necked Stilt *Himantopus mexicanus*
 American Avocet *Recurvirostra americana*

Charadriidae - Lapwings and Plovers

Black-bellied Plover *Pluvialis squatarola*
 Snowy Plover *Charadrius nivosus*
 Semipalmated Plover *Charadrius semipalmatus*
Mountain Plover *Charadrius montanus*
 Killdeer *Charadrius vociferus*

Scolopacidae - Sandpipers, Phalaropes, and Allies

Spotted Sandpiper *Actitis macularius*
Solitary Sandpiper *Tringa solitaria*
 Greater Yellowlegs *Tringa melanoleuca*
 Willet *Tringa semipalmata*
 Lesser Yellowlegs *Tringa flavipes*
 Whimbrel *Numenius phaeopus*
 Long-billed Curlew *Numenius americanus*
 Dunlin *Calidris alpina*
 Least Sandpiper *Calidris minutilla*
 Western Sandpiper *Calidris mauri*
 Short-billed Dowitcher *Limnodromus griseus*
 Long-billed Dowitcher *Limnodromus scolopaceus*
 Wilson's Snipe *Gallinago delicata*
 Wilson's Phalarope *Phalaropus tricolor*
 Red-necked Phalarope *Phalaropus lobatus*

Laridae - Gulls, Terns, and Skimmers

Bonaparte's Gull *Chroicocephalus philadelphia*
Mew Gull *Larus canus*
 Ring-billed Gull *Larus delawarensis*
 California Gull *Larus californicus*
 Herring Gull *Larus argentatus*
Thayer's Gull *Larus thayeri*
Glaucous-winged Gull *Larus glaucescens*
Glaucous Gull *Larus hyperboreus*
 Caspian Tern *Hydroprogne caspia*
 Black Tern *Chlidonias niger*
Common Tern *Sterna hirundo*
 Forster's Tern *Sterna forsteri*

Columbiformes - Pigeons, and Doves

Columbidae - Pigeons and Doves

Rock Pigeon *Columba livia* - I
 Eurasian Collared-Dove *Streptopelia decaocto* - I
 Mourning Dove *Zenaida macroura*

Cuculiformes - Cuckoos and Allies

Cuculidae - Cuckoos, Roadrunners, and Anis

Greater Roadrunner *Geococcyx californianus*

Strigiformes - Owls

Tytonidae - Barn Owls

Barn Owl *Tyto alba*

Strigidae - Typical Owls

Great Horned Owl *Bubo virginianus*

Burrowing Owl *Athene cunicularia*

Caprimulgiformes - Goatsuckers, Oilbirds, and Allies**Caprimulgidae - Goatsuckers**

Lesser Nighthawk *Chordeiles acutipennis*

Apodiformes - Swifts, and Hummingbirds**Apodidae - Swifts**

Vaux's Swift *Chaetura vauxi*

White-throated Swift *Aeronautes saxatalis*

Trochilidae - Hummingbirds

Black-chinned Hummingbird *Archilochus alexandri*

Anna's Hummingbird *Calypte anna*

Rufous Hummingbird *Selasphorus rufus*

Coraciiformes - Rollers, Motmots, Kingfishers, and Allies**Alcedinidae - Kingfishers**

Belted Kingfisher *Megaceryle alcyon*

Piciformes - Puffbirds, Jacamars, Toucans, Woodpeckers, and Allies**Picidae - Woodpeckers and Allies**

Nuttall's Woodpecker *Picoides nuttallii*

Downy Woodpecker *Picoides pubescens*

Northern Flicker *Colaptes auratus*

Falconiformes - Caracaras and Falcons**Falconidae - Caracaras and Falcons**

American Kestrel *Falco sparverius*

Merlin *Falco columbarius*

Peregrine Falcon *Falco peregrinus*

Prairie Falcon *Falco mexicanus*

Passeriformes - Passerine Birds**Tyrannidae - Tyrant Flycatchers**

Olive-sided Flycatcher *Contopus cooperi*

Western Wood-Pewee *Contopus sordidulus*

Willow Flycatcher *Empidonax traillii*

Dusky Flycatcher *Empidonax oberholseri*

Gray Flycatcher *Empidonax wrightii*

Pacific-slope Flycatcher *Empidonax difficilis*

Black Phoebe *Sayornis nigricans*

Eastern Phoebe *Sayornis phoebe*

Say's Phoebe *Sayornis saya*

Vermilion Flycatcher *Pyrocephalus rubinus*

Ash-throated Flycatcher *Myiarchus cinerascens*

Cassin's Kingbird *Tyrannus vociferans*

Western Kingbird *Tyrannus verticalis*

Laniidae - Shrikes

Loggerhead Shrike *Lanius ludovicianus*

Vireonidae - Vireos

Cassin's Vireo *Vireo cassinii*

Warbling Vireo *Vireo gilvus*

Corvidae - Crows and Jays

Western Scrub-Jay *Aphelocoma californica*

American Crow *Corvus brachyrhynchos*

Common Raven *Corvus corax*

Alaudidae - Larks

Horned Lark *Eremophila alpestris*

Hirundinidae - Swallows

Purple Martin *Progne subis*

Tree Swallow *Tachycineta bicolor*

Violet-green Swallow *Tachycineta thalassina*

Northern Rough-winged Swallow *Stelgidopteryx serripennis*

Cliff Swallow *Petrochelidon pyrrhonota*

Barn Swallow *Hirundo rustica*

Aegithalidae - Long-tailed Tits and Bushtits

Bushtit *Psaltriparus minimus*

Troglodytidae - Wrens

Rock Wren *Salpinctes obsoletus*

House Wren *Troglodytes aedon*

Marsh Wren *Cistothorus palustris*

Bewick's Wren *Thryomanes bewickii*

Poliophtilidae - Gnatcatchers and Gnatwrens

Blue-gray Gnatcatcher *Poliophtila caerulea*

Regulidae - Kinglets

Ruby-crowned Kinglet *Regulus calendula*

Turdidae - Thrushes

Western Bluebird *Sialia mexicana*

Mountain Bluebird *Sialia currucoides*

Hermit Thrush *Catharus guttatus*

American Robin *Turdus migratorius*

Mimidae - Mockingbirds and Thrashers

California Thrasher *Toxostoma redivivum*

Sage Thrasher *Oreoscoptes montanus*

Northern Mockingbird *Mimus polyglottos*

Sturnidae - Starlings

European Starling *Sturnus vulgaris* - I

Motacillidae - Wagtails and Pipits

American Pipit *Anthus rubescens*

Bombycillidae - Waxwings

Cedar Waxwing *Bombycilla cedrorum*

Ptiliognatidae - Silky-flycatchers

Phainopepla *Phainopepla nitens*

Calcariidae - Longspurs and Snow Buntings

Chestnut-collared Longspur *Calcarius ornatus*

Parulidae - Wood-Warblers

Orange-crowned Warbler *Oreothlypis celata*

Lucy's Warbler *Oreothlypis luciae*

Virginia's Warbler *Oreothlypis virginiae*

Nashville Warbler *Oreothlypis ruficapilla*

MacGillivray's Warbler *Geothlypis tolmiei*

Common Yellowthroat *Geothlypis trichas*

Yellow Warbler *Setophaga petechia*

Yellow-rumped Warbler *Setophaga coronata*

Black-throated Gray Warbler *Setophaga nigrescens*

Townsend's Warbler *Setophaga townsendi*

Wilson's Warbler *Cardellina pusilla*

Emberizidae - Emberizids

Spotted Towhee *Pipilo maculatus*

California Towhee *Melospiza crissalis*

Chipping Sparrow *Spizella passerina*

Clay-colored Sparrow *Spizella pallida*

Brewer's Sparrow *Spizella breweri*

Vesper Sparrow *Pooecetes gramineus*

Grasshopper Sparrow *Ammodramus savannorum*

Lark Sparrow *Chondestes grammacus*

Black-throated Sparrow *Amphispiza bilineata*

Bell's Sparrow *Artemisiospiza belli canescens*

Savannah Sparrow *Passerculus sandwichensis*

Fox Sparrow *Passerella iliaca*

Song Sparrow *Melospiza melodia*

Lincoln's Sparrow *Melospiza lincolnii*

White-crowned Sparrow *Zonotrichia leucophrys*

Golden-crowned Sparrow *Zonotrichia atricapilla*

Dark-eyed Junco *Junco hyemalis*

Cardinalidae - Cardinals and Allies

Western Tanager *Piranga ludoviciana*

Black-headed Grosbeak *Pheucticus melanocephalus*

Blue Grosbeak *Passerina caerulea*

Lazuli Bunting *Passerina amoena*

Icteridae - Blackbirds

Red-winged Blackbird *Agelaius phoeniceus*

Tricolored Blackbird *Agelaius tricolor*

Western Meadowlark *Sturnella neglecta*

Yellow-headed Blackbird *Xanthocephalus xanthocephalus*

Brewer's Blackbird *Euphagus cyanocephalus*

Great-tailed Grackle *Quiscalus mexicanus*

Brown-headed Cowbird *Molothrus ater*

Hooded Oriole *Icterus cucullatus*

Bullock's Oriole *Icterus bullockii*

Fringillidae - Fringilline and Cardueline Finches and Allies

House Finch *Haemorhous mexicanus*

Purple Finch *Haemorhous purpureus*

Pine Siskin *Spinus pinus*

Lesser Goldfinch *Spinus psaltria*

Lawrence's Goldfinch *Spinus lawrencei*

American Goldfinch *Spinus tristis*

Passeridae - Old World Sparrows

House Sparrow *Passer domesticus* - I